

MAJOR®

NDC 0904-6751-80

ASPIRIN

Low Dose

PAIN RELIEVER (NSAID)
81 mg

Enteric Coated
Safety Coated

Compare to the active
ingredient in Bayer®
Low Dose Aspirin†



1000 Tablets

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)* Pain reliever
*nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains or as recommended by your doctor.
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ take more or for a longer time than directed ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have had stomach ulcers or bleeding problems

■ have 3 or more alcoholic drinks every day while using this product

■ take a blood thinning (anticoagulant) or steroid drug ■ are age 60 or older

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have asthma

■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic

■ you have high blood pressure, heart disease, liver cirrhosis or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for

■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away.

■ you experience any of the following signs of stomach bleeding: ■ vomit blood

■ have bloody or black stools ■ feel faint ■ have stomach pain that does not get better

■ ringing in the ears or a loss of hearing occurs ■ redness or swelling is present

■ pain gets worse or lasts more than 10 days ■ new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison

Control Center (1-800-222-1222) right away.

Directions

■ drink a full glass of water with each dose

■ adults and children 12 years and over: take 4 to 8 tablets every 4

hours not to exceed 48 tablets in 24 hours unless directed by a doctor

■ children under 12 years: do not use unless directed by a doctor

Other information

■ use by expiration date on package

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Purpose

Pain reliever

†This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Bayer® Low Dose Aspirin. 50844 OR6031860016
MAJOR® PHARMACEUTICALS
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152 USA M-17
Re-order No. 700939 Rev. 05/18



Questions or comments? (800) 616-2471