

100 TABLETS 500 MG EACH

Compare to active ingredient in Extra Strength Tylenol®*

ASPIRIN FREE STRONG PAIN & FEVER RELIEF



Acetaminophen EXTRA STRENGTH TABLETS

See New Warnings Information and Directions

MAJOR®

NDC 0904-6730-59

13003-06-18

100 TABLETS 500 MG EACH

Compare to active ingredient in Extra Strength Tylenol®*

ASPIRIN FREE STRONG PAIN & FEVER RELIEF




Acetaminophen EXTRA STRENGTH TABLETS

See New Warnings Information and Directions

MAJOR®

NDC 0904-6730-59

Do not use if carton is open, or if imprinted safety seal under cap is broken or missing.



3 09046 73059 0

Lot #:

Exp. Date:

Drug Facts (continued)
Questions or comments?
Call (800) 616-2471

Distributed By:
MAJOR® PHARMACEUTICALS
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152
Re-Order No. 700921 M-29 Rev. 06/18
*This product is not manufactured or distributed by
McNeil Consumer Healthcare,
owner of the registered trademark Tylenol®

Drug Facts

SAVE CARTON FOR COMPLETE DRUG FACTS

Active ingredient (in each tablet)	Purpose
Acetaminophen 500 mg	Pain reliever/ Fever reducer

Uses

Temporarily reduces fever and relieves minor aches and pains due to:

- headache • muscular aches • backache • minor pain of arthritis • common cold • toothache • premenstrual and menstrual cramps

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are allergic to acetaminophen or any of the inactive ingredients in this product

Drug Facts (continued)

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away, (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not take more than directed (see overdose warning)
adults and children 12 years and over:

- take 2 tablets every 6 hours while symptoms last
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor

children under 12 years: ask a doctor

Other information store at room temperature

Inactive ingredients Povidone, pregelatinized starch, sodium starch glycolate, stearic acid