

MAJOR®

Fast, Long Lasting Relief

Sore Throat Lozenges

CHERRY

NDC 0904-6255-49

MAJOR®

Fast, Long Lasting Relief

Sore Throat Lozenges

Menthol/Benzocaine

For Temporary Relief of Minor Sore Throat Pain

Oral Anesthetic

18 Lozenges

CHERRY

Compare to the active ingredients in Cepacol®

PN-320232



Distributed by MAJOR® PHARMACEUTICALS 17177 N LAUREL PARK DRIVE, SUITE 233, LITTLETON, CO 80120 USA M-130 Re-order No. 700608 Rev. 8/18 3 09046 25549 9

Gluten-Free

No Ink or Coating Lot/Exp Info

WARNING: TAMPER-EVIDENT PACKAGING. DO NOT USE IF OUTER BOX OR BLISTER IS OPEN OR DAMAGED. This product is not manufactured or distributed by Reckitt Benckiser owner of the registered trademark Cepacol®.

Drug Facts (continued) Inactive ingredients: Corn starch, FD&C Red #40, glucose syrup, glycerin, medium chain triglycerides, natural & artificial flavors, propylene glycol, soybean oil, sucrose and water. Questions? Call 1-800-616-2471, Monday through Friday, 9AM - 4PM EST

Drug Facts Active ingredients (in each lozenge) Benzocaine 15 mg, Menthol 3.6 mg, Oral anesthetic Purpose Uses temporarily relieves these symptoms: occasional minor irritation, pain, sore mouth and sore throat. Warnings Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: pale, gray, or blue colored skin (cyanosis), headache, rapid heart rate, shortness of breath, dizziness or lightheadedness, fatigue or lack of energy. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly. Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics. Do not use for teething, in children under 2 years of age. Stop use and ask doctor if sore throat is severe or irritation, pain or redness persists or worsens, sore mouth symptoms do not improve in 7 days. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away. Directions adults and children allow 1 lozenge to dissolve slowly or as directed by a doctor every two hours as necessary or as directed by a doctor. children under 5 years do not use. Other information store at room temperature protect from moisture

No Ink or Coating

No Ink or Coating

No Ink or Coating